



Short Communication

Role of pigeons in the spread of Newcastle disease

M. Geetha*¹, L. Gunaseelan², P. I. Ganesan¹ and K. Kumanan³

¹Department of Veterinary Epidemiology and Preventive Medicine, ²Department of Veterinary Public Health and Epidemiology, Madras Veterinary College, Chennai, Tamil Nadu, India; ³Tamil Nadu Veterinary and Animal Sciences University, Chennai, Tamil Nadu, India

Abstract

Newcastle disease (ND) is one of the major viral diseases of poultry causing great economic losses to the poultry industry. This study was aimed to assess the role of pigeons in the epidemiology of ND. A total of 70 cloacal swabs/droppings from pigeons were collected and inoculated into specific pathogen free embryonated chicken eggs for isolation of Newcastle disease virus (NDV). Only one NDV isolate was obtained out of 70 cloacal swabs inoculated and the isolate was characterized as velogenic pathotype based on mean death time (MDT) and intracerebral pathogenicity index (ICPI). It was concluded that pigeons may play an important role in the transmission of NDV to domestic chicken. Hence biosecurity measures have to be strictly followed for avoiding the contact between free living pigeons and domestic chicken for effective prevention of ND.

Keywords: Newcastle disease; pigeon; biosecurity

To cite this article: Geetha M, L Gunaseelan, PI Ganesan and K Kumanan, 2014. Role of pigeons in the spread of Newcastle disease. *Res. Opin. Anim. Vet. Sci.*, 4(4): 170-171.

Introduction

Newcastle disease remains a constant threat to the poultry industry and is a limiting disease for poultry producers worldwide (Cattoli et al., 2011). It may represent a bigger drain on the world economy than any other animal viral disease (Alexander, 2003). Apart from commercial poultry, a wide range of captive and free living birds are susceptible and can act as primary source of ND infection to chicken (Kouwenhoven, 1993; Alexander and Senne, 2008). Limited work has been published to know the role of free living birds like Desi chicken, caged pet birds, turkeys, geese, pigeons, sparrows, crows etc. in the spread of the ND to commercial chicken (Raghavan et al., 1998). Hence, this study was aimed to assess the role of pigeons in the spread of ND.

Materials and Methods

A total number of 70 cloacal swabs/droppings from free living pigeons were collected in phosphate

buffered saline at Chennai, Tamil Nadu, India, and centrifuged at 1500g for 15 min at 4°C. The supernatant was treated with penicillin at the rate of 10,000 IU/ml and streptomycin at the rate of 10 mg/ml used for virus isolation. Virus isolation, identification and characterization were carried out by the procedures described by Alexander and Senne (2008). The supernatants of cloacal swabs were inoculated into 9-10 day old specific pathogen free embryonated chicken eggs through allantoic cavity route and incubated at 37°C. Those embryos died within 24 hours after inoculation was considered as non-specific and those embryos that died after 24 hours were chilled. Amnio-allantoic fluid (AAF) collected from the dead embryos were subjected to haemagglutination (HA) test with 1 percent washed chicken erythrocytes and all the dead embryos were examined for the presence of characteristic NDV lesions. The AAF samples which were not agglutinating chicken erythrocytes at first passage were subjected to two more blind passages in embryonated eggs. Spot HA test was carried out at each passage level and the samples which showed the HA

Corresponding author: M. Geetha, Department of Veterinary Epidemiology and Preventive Medicine, Madras Veterinary College, Chennai, Tamil Nadu, India

activity were confirmed by haemagglutination inhibition (HI) test by using specific NDV antiserum raised at the Department of Animal Biotechnology, Madras Veterinary College, Chennai. The viral isolate obtained was characterized by mean death time (MDT) in specific pathogen free embryonated chicken eggs and intracerebral pathogenicity index (ICPI) in day old chicks.

Results and Discussion

Out of the 70 number of cloacal swabs from pigeons inoculated into the embryonated chicken eggs, only one sample yielded NDV isolate at III passage level. The embryo death was noticed at 48 hours post inoculation and the HA titre of AAF was found to be 64. The HI titre reciprocal of the serum dilution was 32. The dead embryo showed hemorrhage on the cranium. The viral isolate has ICPI value of 1.975 and as per Werner et al. (1999), NDV with ICPI value of above 1 are considered as velogenic. Mean death time of the NDV isolate obtained in this study was 72 hours and identified as mesogenic pathotype based on this test. Eventhough MDT was considered as an important tool in characterizing different isolates, it has been reported to be imprecise particularly when used to characterize isolates from hosts other than chicken (Alexander, 1988). Based on the ICPI value the NDV isolate obtained in this study might be of velogenic pathotype.

Newcastle disease virus obtained from pigeons is an indication that the free living pigeons can be a natural reservoir of NDV and act as a source of infection to commercial poultry. Velogenic NDV was isolated from pigeons by Pearson et al. (1987) and Meulemans et al. (2002). An earlier report by Alexander and Parsons (1986) indicated increase of virulence of NDV isolates from pigeons. Such possibilities caution the importance of free living pigeons as NDV reservoirs. Identification of NDV isolates in this study is evidence of the need for continuous characterization of NDV strains of all pathotypes which will provide a better understanding of the diversity that exists as suggested by King and Seal (1997). Single stranded RNA viruses which lack of proof reading and postreplicative error correction mechanisms are expected to have high mutation rate and, therefore, to evolve rapidly (Koonin and Dolja, 1993). Immune system of the birds may force the virus to evolve more rapidly in order to create escape mutants, which in turn might lead to the emergence of a few mutants and avoid clearance by the hosts immune system and such mutants have the ability to spread widely and cause epidemics in chickens (Kee t al., 2001).

Conclusion

Hence in this scenario, biosecurity measures should be strictly adhered to break the contact of commercial poultry with pigeons for prevention of spread of NDV from pigeons to commercial poultry.

References

- Alexander, D.J. and Parsons, G. 1986. Protection of chickens against challenge with the variant virus responsible for Newcastle disease in 1984 by conventional vaccination. *Veterinary Record*, 118: 176-177.
- Alexander, D.J. 1988. Newcastle disease, 1st edition, Kluwer Academic Publishers, London. Pp: 147-183.
- Alexander, D.J. 2003. Newcastle disease and other avian paramyxoviruses and pneumoviruses. In: Saif, Y.M. (eds), Diseases of poultry, Ames: Iowa State University Press, Blackwell, Oxford, UK, pp: 75-116.
- Alexander, D.J. and Senne, D.A. 2008. Newcastle disease and other avian paramyxoviruses and pneumoviruses, in: Saif, Y.M. (eds), Diseases of poultry, Ames: Iowa State University Press, Blackwell, Oxford, UK, Pp: 75-116.
- Cattoli, G., Susta, L., Terregino, C. and Brown, C. 2011. Newcastle disease: a review of field recognition and current methods of laboratory detection. *Journal of Veterinary Diagnostic Investigation*, 23: 637-656.
- Ke, G.M., Liu, H.J., Lin, M.Y., Chen, J.H. Tsai, S.S. and Chang, P.C. 2001. Analysis of sequence and haemagglutinin activity of the HN glycoprotein of Newcastle disease virus. *Journal of virological Methods*, 97: 1-11.
- King, D.J. and Seal, B.S. 1997. Biological and molecular characterization of Newcastle disease viral isolates from surveillance of live bird markets in the North-Eastern United States. *Avian Diseases*, 41: 683-689.
- Koonin, E.V. and Dolja, V.V. 1993. Evolution and taxonomy of positive-strand RNA viruses: implications of comparative analysis of amino acid sequences. *Critical Review in Biochemistry and Molecular Biology*, 28: 375-430.
- Kouwenhoven, B. 1993. Newcastle disease. In: McFerren, J.B. and McNulty, M.S. (ed), Virus Infection of Birds. Elsevier science publishers, Amsterdam and New York. Pp: 341-361.
- Meulemans, G., Van den Berg, T.P., Decaesstecker, M. and Boschmans, M. 2002. Evolution of pigeon Newcastle disease virus strains. *Avian Pathology*, 31: 515-519.
- Pearson, J.E., Senne, D.A., Alexander, D.J., Taylor, W.D., Peterson, L.A. and Russell, P.H. 1987. Characterization of Newcastle disease virus (avian paramyxovirus-1) isolated from pigeons, *Avian Diseases*, 31: 105-111.
- Raghavan, V.S., Kumanan, K., Thirumurugan, G. and Nachimuthu, K. 1998. Comparison of various diagnostic methods in characterizing Newcastle disease virus isolates from desi chickens, *Tropical Animal Health Production*, 30: 287-293.
- Werner, O., Romer-Oderdorfer, A., Kollner, B., Manvell, R.J. and Alexander, D.J. 1999. Characterization of avian paramyxovirus type 1 strains isolated in Germany during 1992-1996. *Avian Pathology*, 28: 79-88.